

Second-parameter Globular Clusters in the Milky Way and in M33 as Tracers of Mass Loss from M31 in the Early Epoch?

Valery V. Kravtsov

Sternberg Astron. Ins., 13, University Ave., Moscow 119899, Russia

Abstract. I suggest that the bulk of the "young", second-parameter globular clusters (SPGCs) observed in the outer halo of our Galaxy and recently found in other massive spiral of the Local Group (LG), M33, may have originated due to mass outflow from M31 and subsequent accretion of gas on the Galaxy and M33 in the early epoch.

1. Problem and Main Questions

There are contradictory conclusions about the dwarf spheroidal (dSph) galaxies' tidal dissolution and its rate, as well as a number of observational constraints on this process in the Galaxy (e.g., van den Bergh 1996; Gilmore et al. 1999; Shetrone et al. 2000). Moreover, preliminary results on the CMD study of GCs in massive spirals of LG suggest a fraction of SPGCs to be at least low among GCs in M31 (Rich et al. 2001) and to be high as for M33 GC population (Sarajedini et al. 1998). Hence there appear doubts about tidal disruption/stripping of dSphs as universal mechanism of and main contributor to a formation of populations of the SPGCs in the above three galaxies. Note also that Chandar et al. (2001) have estimated the total number of GCs in M 33 to be 75 ± 14 which gives a specific frequency significantly higher than in other late-type spirals.

By now, it is established that the Galactic population of SPGCs shows a number of basic distinctive characteristics as compared with those of other GCs in the halo of the Milky Way (e.g., Da Costa 1994; van den Bergh 1996). Some of these characteristics imply that the Galactic SPGCs are very likely to have an accretive nature. However, if a tidal dissolution of dSphs is not main and the only contributor to the formation of population(s) of the SPGCs in our Galaxy (and in M33) then the following vexed questions arise.

What is a material accreted and where did it originate and come from? Why has the formation of SPGCs lasted after formation of the bulk of halo objects in rapid collapse?

2. Putative Role of the Mass Outflows

The high-redshift Lyman break galaxies (LBGs) are suggested to be progenitors of the present-day massive spheroids being in particular components of luminous early-type spirals (Friaca & Terlevich 1999). As summarized by Heckman (2000) starburst-driven galactic superwinds observed in LBGs play important role in

the mass outflow from them. During main, most powerful episode (~ 2 Gyr) of formation of the Pop. II stars in a star forming galaxy like M31, its outflows may carry off the galaxy the total mass of gas as large as mass of the stellar bulge and spheroid, with typical speeds ranging from **a few 10^2 to 10^3 km/s** (Pettini et al. 2001). Some a few tens percent of this mass ($\Delta M \sim 10^{11} M_\odot$ or so) with velocities exceeding the escape velocity of M31 could have quited the galaxy, portion of which might be accreted on our Galaxy (and M33). If we accept present distance between the Galaxy and M31 (~ 0.7 Mpc), ratio of their mass ($M_G/M_A \sim 1/1.5$), and also a mean velocity of expanding gas to be of order **400 km/s**, then we easily find that a time spent by gas to reach a region of the Lagrangian point between the two galaxies, may be as long as ~ 1 Gyr. **An upper limit of mass δM of gas accreted on the Galaxy** is estimated from a formula obtained (in the approach of gas accretion in gravity potential $\varphi \sim R^{-1}$ of a field of less massive companion in a binary system) and kindly offered by Postnov (2000, private communication):

$$\delta M \sim \frac{1}{4} \left(\frac{M_G}{M_A} \right)^2 \Delta M \sim \frac{1}{10} \Delta M \sim 10^{10} M_\odot$$

Even if a mass of really accreted gas was an order of magnitude lower than the upper limit estimated and if in turn approximately ten percent of this mass was converted into objects, i.e., $\sim 10^8 M_\odot$, that would be equal to a **total mass** of the bulk of **SPGCs**, of a few **dSphs**, and of some portion of **field stars** which may have formed due to such a process in the outer halo of our Galaxy.

Similarly, it is possible to conclude that objects assumed to form from a gas accreted onto M33 might comprise a total mass of order **a few $10^6 M_\odot$** . Some **10 to 20 GCs** (with fraction of **field stars** formed) with cluster's typical mass of order $10^5 M_\odot$ may amount just to this mass.

Acknowledgments. I am deeply thankful for the IAU travel grant. Also, I am very grateful to Dr. A.A. Suchkov for his encouraging and valuable comments on the scenario presented, and to Drs. K. Postnov, Yu.A. Shchekinov, Yu.N. Efremov, N.I. Shakura, and M.V. Sazhin for helpful discussions.

References

- Chandar, R., Bianchi, L., & Ford, H. C. 2001, A&A, 366, 498
 Da Costa, G. S. 1994, In ESO Conference and Workshop Proc., 51, The Local Group: Comparative and Global Properties, ed. A. Layden, R.C. Smith & J. Strom, 101
 Friaca, A. C. S. & Terlevich, R. J. 1999, MNRAS, 305, 90
 Gilmore, G., Hernadez, X., Valls-Gabaud, D. 1999, astro-ph/9910409
 Heckman, T. M. 2000, astro-ph/0009075
 Pettini, M., Steidel, C. C., Shapley, A. E., et al. 2001, astro-ph/0101254
 Rich, R. M., Corsi, C. E., Cacciari, C. et al. 2001, this volume
 Sarajedini, A., Geisler, D., Harding, P., & Schommer, R. 1998, ApJ, 508, L37
 Shetrone, M., Cote, P., Sargent, W. L. W. 2000, astro-ph/0009505
 van den Bergh, S. 1996, PASP, 108, 986